

## **Neospora caninum and neosporosis**

Patrick S. Craig ASc, DVM, PhD (awarded)

*Neospora caninum* is a protozoan cyst forming apicomplexan parasite that causes neosporosis, notably in cattle (*Bos taurus*) and domestic dogs (*Canis familiaris*) [1,2]. It is closely related to *Toxoplasma gondii* and has emerged as a major cause of reproductive failure in cattle worldwide [2]. The parasite was first described in a dog with encephalomyelitis and myositis [1] and was later described in calves with myeloencephalitis [3,4]. *Neospora caninum* has been isolated from a variety of animal host species, such as the dog (Nc1;[5] and Nc-Liverpool [6]), cattle (BPA1 and BPA2 [7]), sheep (NC-Sheep [8]) and water buffalo (NCBrbuf-1,2,3,4,5 [9]). Antibodies to *N. caninum* have also been identified in the captive sika deer (*Cervus Nippon*), mouflon (*Ovis musimon*), fallow deer (*Dama dama*), moose (*Alces alces*), European bison (*Bison bonasusbonasus* L.), wild rabbits (*Oryctolagus cuniculus*), brown hares (*Lepus europeus*) and dolphin (*Tursiops truncatus*) (reviewed in [10]). Apart from the dog (*Canis familiaris*), other canids have also been considered as potential definitive hosts of *N. caninum*. To date, coyotes [11], wolves (*Canis lupus*) [12] and dingoes [13] have also been named as definitive hosts because they have been shown to shed oocysts after being fed infected tissues, while many other domestic and wild mammal species have been identified as intermediate hosts [10,14].

Horses are an intermediate hosts of *Neospora hughesi* [15], which was identified in an aborted equine foetus [16]. The parasite was later isolated from an 11-year old horse and described as a new species, with molecular differences to *N. caninum* [15]. It is still unknown whether *N. hughesi* is the sole species of *Neospora* that infects horses or if *N. caninum* also plays a part in infection as both species can cross react serologically with each other [17].

The parasite has three asexual infectious development stages, i.e. tachyzoites, bradyzoites and sporozoites [18]. The sexual stages of *Neospora* have not been described, but it is likely that the schizont (asexual) and gamont (sexual) stages also exist in the gut epithelium of the definitive hosts as was shown for *T. gondii* when kittens were fed *Toxoplasma* cysts, since these two parasites are closed related [19]. *Neospora caninum* is regarded as one of the most important infectious causes of abortion in cattle worldwide, yet there are no definitive studies that quantify losses due to neosporosis in the cattle industry; however, losses are estimated to be in the

millions of dollars per year [20,21]. The seroprevalence of *N. caninum* was found to be approximately 26% (PhD thesis, Patrick Craig) and has been estimated at approximately 12.9% in the UK [22], whereas it was as low as 2.8% in Sweden [23] and reached 55.9% in Romania [24], 14-40% in the Americas [25], 6-36% in Asia [26] and 6-21% in Oceania [27].

The two major routes of *N. caninum* transmission are horizontal, where cattle ingest sporulated *N. caninum* oocysts and vertical transmission, which includes transmission of the agent to the foetus during pregnancy, either following reactivation of bradyzoites in the infected dam or *de novo* infection of the dam during pregnancy [28]. Vertical transmission contributes significantly to the persistence of *N. caninum* in a herd by propagating the infection to successive generations. The prevalence of congenital infection varies with reported infection rates ranging from approximately 40% to 95% [29-33]. The shedding of oocysts by infected canids in cattle-feeding areas is a likely cause of horizontal transmission that could play an important role in infected herds. Cow-to-cow transmission has not been observed to date, but many authors have looked at the possibility of infection via contaminated semen from infected bulls [34-37]. However, artificial insemination of cows with semen *in vitro* contaminated with *N. caninum* tachyzoites ( $6.5 \times 10^7$  and  $1.8 \times 10^7$  on day one and two, respectively) failed to induce infection [34]. Although one animal developed a low antibody titre of 1:80 in the direct agglutination test at day ten after insemination, it was negative after 45 days, which shows that the parasites were able to stimulate the immune system of this animal without causing infection.

The pathogenesis of *N. caninum* induced abortion in cattle is yet to be fully elucidated. *Neospora caninum* infection in cows is mainly manifested in the placenta and foetal tissues following a maternal parasitaemia. Experimental studies have shown that infection of pregnant cows in early gestation leads to foetal death, which could be due to both extensive placental necrosis and necrosis in foetal tissues, such as the liver and brain [38-44]. Changes observed in first trimester fetuses include necrosis and apoptosis in the CNS (brain and spinal cord), liver, lung, kidney, spleen and thymus. The parasite seems to replicate in the heart without causing cell death (studied extensively in PhD thesis, Chapter 3). In fetuses following infection of the dams in mid to late gestation, mild to moderate histological changes are detected in foetal tissues and are restricted mainly to the CNS [38].

The host immune response towards *N. caninum* is of paramount importance in controlling parasite multiplication. *Neospora caninum* is an obligate intracellular pathogen, which means that cell mediated immune responses are likely to play a pivotal role in protective immunity, by reducing the multiplication of parasites within the host, hence reducing parasitaemia [45,46]. Protective immunity to intracellular pathogens is associated with type 1 T helper cells (Th1 cells), which secrete pro-inflammatory cytokines, such as interferon gamma (IFN- $\gamma$ ), tumour necrosis factor alpha (TNF- $\alpha$ ) and interleukin (IL)12 [47]. The Th2 type anti-inflammatory cytokines, namely IL-4, and regulatory cytokines which includes IL-10 and transforming growth factor beta (TGF- $\beta$ ) are produced at the foeto-maternal interface in the placenta and can counteract the effect of the pro-inflammatory cytokines [48]. The Th2 cytokines are associated with implantation of the foetus and maintenance of early pregnancy by suppression of local inflammatory responses [49,50].


For the diagnosis of bovine neosporosis, clinical history, epidemiological data, information about the abortion pattern and foetal age are important factors that should be considered [51]. The definitive diagnosis of neosporosis can be very difficult, because infection does not always result in abortion and even demonstration of *N. caninum* infection, both histologically and immunohistologically, does not give conclusive evidence that the parasite is the cause of the abortion [52]. With regards to post-mortem diagnosis in aborted foetuses, the ideal diagnostic samples include both the aborted foetus and the placenta, together with sera from the dam. If this is not feasible, samples from brain, heart and liver of the foetus should be submitted [51]. The brain is the most reliable tissue for the diagnosis, but the probability of diagnosing the infection increases when other tissues, such as heart and liver, are analysed [51]. Histology has been the most commonly used method for parasite detection initially, but immunohistological detection of parasites in foetal brain, lung, liver and heart has been used to confirm the presence of the agent [53,54]. The PCR technique also plays an important role in the diagnosis of *N. caninum* infection when used in aborted foetal tissues [55-58] and other samples such as amniotic fluid [59] and cerebrospinal fluid [60,61]. The advantage of the PCR is its high specificity and high sensitivity and therefore the ability to detect small amounts of *N. caninum* DNA in a large quantity of tissue; PCR also works well when foetal tissues are autolysed, which is often the case with *Neospora* abortions [51]. Serological tests have the advantage that they can be applied *intra*

*vitam* and are suitable techniques for processing large numbers of samples [62]. There is a variety of serological assays available for the detection of *N. caninum* antibodies in cattle. The immunofluorescent antibody test (IFAT) has been widely used to detect *N. caninum* specific antibody in maternal serum or foetal fluids. In addition to the IFAT, a number of *N. caninum* specific enzyme-linked immunosorbent assays (ELISAs) have been described, which utilise either whole fixed *N. caninum* tachyzoites, aqueous or detergent-soluble tachyzoite extracts, tachyzoite antigens incorporated into immunostimulating complexes (iscoms) or recombinant tachyzoite antigens [63-67].

Seropositivity to *Neospora caninum* in cattle of different age groups from three different locations in Jamaica (sampled in 2012)

Age groups	Number of animals	Positives (%)	Negatives	Pregnant (%) <sup>a</sup>
<b>0-2</b>	16	5 (31.3)	11	4 (25.0)
<b>3-5</b>	229	53 (23.1)	176	43 (18.8)
<b>6-8</b>	151	33 (21.9)	118	26 (17.2)
<b>9-11</b>	66	22 (33.3)	44	10 (15.1)
<b>&gt;12</b>	30	13 (43.3)	17	4 (13.3)
<b>unknown</b>	7	2 (28.6)	5	0 (0.0)

<sup>a</sup> overall number of pregnant cows sampled



Ultrastructural features of *Neospora caninum*-infected human HepG2 cells cultured in vitro with 95% O<sub>2</sub> and 5% CO<sub>2</sub>. **A.** *Neospora caninum* tachyzoites (T) are present in hepatocytes with numerous mitochondria (M) that are predominantly located in close contact with the parasitophorous vacuole membrane. **B.** Hepatocyte with two parasitophorous vacuoles filled with tachyzoites (T). Numerous mitochondria (M) are located in the perinuclear region and in association with the PVM. Lead citrate/uranyl acetate staining, FIB-SEM Nanotomography. (This photomicrograph is the sole property of Patrick Craig and the University of Liverpool).

## References

1. Bjerkas I, Mohn SF, Presthus J (1984) Unidentified cyst-forming Sporozoon causing encephalomyelitis and myositis in dogs. *Zeitschrift fur Parasitenkunde* 70: 271-274.
2. Dubey JP, Lindsay DS (1996) A review of *Neospora caninum* and neosporosis. *Veterinary Parasitology* 67: 1-59.
3. Parish SM, Maag-Miller L, Besser TE, Weidner JP, McElwain T, et al. (1987) Myelitis associated with protozoal infection in newborn calves. *Journal of the American Veterinary Medical Association* 191: 1599-1600.
4. O'Toole D, Jeffrey M (1987) Congenital sporozoan encephalomyelitis in a calf. *Veterinary Record* 121: 563-566.

5. Dubey JP, Hattel AL, Lindsay DS, Topper MJ (1988) Neonatal *Neospora caninum* infection in dogs: isolation of the causative agent and experimental transmission. *Journal of the American Veterinary Medical Association* 193: 1259-1263.
6. Barber J, Trees AJ, Owen M, Tennant B (1993) Isolation of *Neospora caninum* from a British dog. *Veterinary Record* 133: 531-532.
7. Conrad PA, Barr BC, Sverlow KW, Anderson M, Daft B, et al. (1993) In vitro isolation and characterization of a *Neospora* sp. from aborted bovine fetuses. *Parasitology* 106: 239-249.
8. Koyama T, Kobayashi Y, Omata Y, Yamada M, Furuoka H, et al. (2001) Isolation of *Neospora caninum* from the brain of a pregnant sheep. *Journal of Parasitology* 87: 1486-1488.
9. Rodrigues AAR, Gennari SM, Aguiar DM, Sreekumar C, Hill DE, et al. (2004) Shedding of *Neospora caninum* oocysts by dogs fed tissues from naturally infected water buffaloes (*Bubalus bubalis*) from Brazil. *Veterinary Parasitology* 124: 139-150.
10. Dubey JP, Schares G, Ortega-Mora LM (2007) Epidemiology and control of neosporosis and *Neospora caninum*. *Clinical Microbiology Reviews* 20: 323-367.
11. Gondim LFP, McAllister MM, Pitt WC, Zemlicka DE (2004) Coyotes (*Canis latrans*) are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* 34: 159-161.
12. Dubey JP, Jenkins MC, Rajendran C, Miska K, Ferreira LR, et al. (2011) Gray wolf (*Canis lupus*) is a natural definitive host for *Neospora caninum*. *Veterinary Parasitology* 181: 382-387.
13. King JS, Šlapeta J, Jenkins DJ, Al-Qassab SE, Ellis JT, et al. (2010) Australian dingoes are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* 40: 945-950.
14. Dubey JP, Schares G (2011) Neosporosis in animals-The last five years. *Veterinary Parasitology* 180: 90-108.
15. Marsh AE, Barr BC, Packham AE, Conrad PA (1998) Description of a new *Neospora* species (Protozoa: Apicomplexa: Sarcocystidae). *Journal of Parasitology* 84: 983-991.
16. Dubey JP, Porterfield ML (1990) *Neospora caninum* (Apicomplexa) in an aborted equine fetus. *Journal of Parasitology* 76: 732-734.
17. Gondim LFP, Lindsay DS, McAllister MM (2009) Canine and bovine neospora caninum control sera examined for cross-reactivity using neospora caninum and neospora hughesi indirect fluorescent antibody tests. *Journal of Parasitology* 95: 86-88.
18. Dubey JP (1999) Recent advances in *Neospora* and neosporosis. *Veterinary Parasitology* 84: 349-367.
19. Frenkel JK, Dubey JP, Miller NL (1970) *Toxoplasma gondii* in cats: Fecal stages identified as coccidian oocysts. *Science* 167: 893-896.
20. Dubey JP (2003) Review of *Neospora caninum* and neosporosis in animals. *The Korean journal of parasitology* 41: 1-16.
21. Reichel MP, Alejandra Ayanegui-Alcérreca M, Gondim LFP, Ellis JT (2013) What is the global economic impact of *Neospora caninum* in cattle - The billion dollar question. *International Journal for Parasitology* 43: 133-142.
22. Woodbine KA, Medley GF, Moore SJ, Ramirez-Villaescusa A, Mason S, et al. (2008) A four year longitudinal sero-epidemiology study of *Neospora caninum* in adult cattle from 114 cattle herds in south west England: Associations with age, herd and dam-offspring Pairs. *BMC Vet Res* 4.
23. Loobuyck M, Frössling J, Lindberg A, Björkman C (2009) Seroprevalence and spatial distribution of *Neospora caninum* in a population of beef cattle. *Preventive Veterinary Medicine* 92: 116-122.
24. Gavrea RR, Cozma V (2010) Seroprevalence of *Neospora caninum* in cows with reproductive failure in center and northwest of Romania. *Sci Parasitol* 11: 67-70.
25. Moore DP (2005) Neosporosis in South America. *Veterinary Parasitology* 127: 87-97.
26. Koiwai M, Hamaoka T, Haritani M, Shimizu S, Kimura K, et al. (2005) Proportion of abortions due to neosporosis among dairy cattle in Japan. *Journal of Veterinary Medical Science* 67: 1173-1175.
27. Hall CA, Reichel MP, Ellis JT (2006) Prevalence of *Neospora caninum* infection in Australian (NSW) dairy cattle estimated by a newly validated ELISA for milk. *Veterinary Parasitology* 142: 173-178.

28. Trees AJ, Williams DJL (2005) Endogenous and exogenous transplacental infection in *Neospora caninum* and *Toxoplasma gondii*. *Trends in Parasitology* 21: 558-561.
29. Pan Y, Jansen GB, Duffield TF, Hietala S, Kelton D, et al. (2004) Genetic susceptibility to *Neospora caninum* infection in Holstein cattle in Ontario. *Journal of dairy science* 87: 3967-3975.
30. Bergeron N, Fecteau G, Paré J, Martineau R, Villeneuve A (2000) Vertical and horizontal transmission of *Neospora caninum* in dairy herds in Québec. *Canadian Veterinary Journal* 41: 464-469.
31. Romero JJ, Frankena K (2003) The effect of the dam-calf relationship on serostatus to *Neospora caninum* on 20 Costa Rican dairy farms. *Veterinary Parasitology* 114: 159-171.
32. Davison HC, French NP, Trees AJ (1999) Herd-specific and age-specific seroprevalence of *Neospora caninum* in 14 British dairy herds. *Veterinary Record* 144: 547-550.
33. Björkman C, McAllister MM, Frössling J, Näslund K, Leung F, et al. (2003) Application of the *Neospora caninum* IgG avidity ELISA in assessment of chronic reproductive losses after an outbreak of neosporosis in a herd of beef cattle. *Journal of Veterinary Diagnostic Investigation* 15: 3-7.
34. Canada N, Meireles CS, Ferreira P, Correia da Costa JM, Rocha A (2006) Artificial insemination of cows with semen in vitro contaminated with *Neospora caninum* tachyzoites failed to induce neosporosis. *Veterinary Parasitology* 139: 109-114.
35. Ortega-Mora LM, Ferre I, Del-Pozo I, Caetano-Da-Silva A, Collantes-Fernández E, et al. (2003) Detection of *Neospora caninum* in semen of bulls. *Veterinary Parasitology* 117: 301-308.
36. Serrano-Martínez E, Ferre I, Martínez A, Osoro K, Mateos-Sanz A, et al. (2007) Experimental neosporosis in bulls: Parasite detection in semen and blood and specific antibody and interferon-gamma responses. *Theriogenology* 67: 1175-1184.
37. Jozani RJ, Asadpour R, Nematollahi A, Hosseininejad M (2012) Detection of non-spermatozoal cells of *Neospora caninum* in fresh semen of naturally infected bulls. *Acta Scientiae Veterinariae* 40.
38. Gibney EH, Kipar A, Rosbottom A, Guy CS, Smith RF, et al. (2008) The extent of parasite-associated necrosis in the placenta and foetal tissues of cattle following *Neospora caninum* infection in early and late gestation correlates with foetal death. *International Journal for Parasitology* 38: 579-588.
39. Maley SW, Buxton D, Macaldowie CN, Anderson IE, Wright SE, et al. (2006) Characterization of the Immune Response in the Placenta of Cattle Experimentally Infected with *Neospora caninum* in Early Gestation. *Journal of Comparative Pathology* 135: 130-141.
40. Macaldowie C, Maley SW, Wright S, Bartley P, Esteban-Redondo I, et al. (2004) Placental pathology associated with fetal death in cattle inoculated with *Neospora caninum* by two different routes in early pregnancy. *Journal of Comparative Pathology* 131: 142-156.
41. Rosbottom A, Gibney EH, Guy CS, Kipar A, Smith RF, et al. (2008) Upregulation of cytokines is detected in the placentas of cattle infected with *Neospora caninum* and is more marked early in gestation when fetal death is observed. *Infection and Immunity* 76: 2352-2361.
42. Rosbottom A, Guy CS, Gibney EH, Smith RF, Valarcher JF, et al. (2007) Peripheral immune responses in pregnant cattle following *Neospora caninum* infection. *Parasite Immunology* 29: 219-228.
43. Williams DJL, Guy CS, McGarry JW, Guy F, Tasker L, et al. (2000) *Neospora caninum*-associated abortion in cattle: The time of experimentally-induced parasitaemia during gestation determines foetal survival. *Parasitology* 121: 347-358.
44. Rosbottom A, Gibney H, Kaiser P, Hartley C, Smith RF, et al. (2011) Up regulation of the maternal immune response in the placenta of cattle naturally infected with *Neospora caninum*. *PLoS ONE* 6.
45. Innes EA, Wright S, Bartley P, Maley S, Macaldowie C, et al. (2005) The host-parasite relationship in bovine neosporosis. *Veterinary Immunology and Immunopathology* 108: 29-36.
46. Almeria S, Araujo RN, Darwich L, Dubey JP, Gasbarre LC (2011) Cytokine gene expression at the materno-foetal interface after experimental *Neospora caninum* infection of heifers at 110 days of gestation. *Parasite Immunology* 33: 517-523.
47. Baszler TV, Long MT, McElwain TF, Mathison BA (1999) Interferon- $\gamma$  and interleukin-12 mediate protection to acute *Neospora caninum* infection in BALB/c mice. *International Journal for Parasitology* 29: 1635-1646.

48. Entrican G (2002) Immune regulation during pregnancy and host-pathogen interactions in infectious abortion. *Journal of Comparative Pathology* 126: 79-94.
49. Chaouat G, Zourbas S, Ostojic S, Lappree-Delage G, Dubanchet S, et al. (2002) A brief review of recent data on some cytokine expressions at the materno-foetal interface which might challenge the classical Th1/Th2 dichotomy. *Journal of Reproductive Immunology* 53: 241-256.
50. Wegmann TG, Lin H, Guilbert L, Mosmann TR (1993) Bidirectional cytokine interactions in the maternal-fetal relationship: Is successful pregnancy a TH2 phenomenon? *Immunology Today* 14: 353-356.
51. Ortega-Mora LM, Fernández-García A, Gómez-Bautista M (2006) Diagnosis of bovine neosporosis: Recent advances and perspectives. *Acta Parasitologica* 51: 1-14.
52. Thurmond MC, Hietala SK, Blanchard PC (1999) Predictive values of fetal histopathology and immunoperoxidase staining in diagnosing bovine abortion caused by *Neospora caninum* in a dairy herd. *Journal of Veterinary Diagnostic Investigation* 11: 90-94.
53. Lindsay DS, Dubey JP (1989) Immunohistochemical diagnosis of *Neospora caninum* in tissue sections. *American Journal of Veterinary Research* 50: 1981-1983.
54. Boger LA, Hattel AL (2003) Additional evaluation of undiagnosed bovine abortion cases may reveal fetal neosporosis. *Veterinary Parasitology* 113: 1-6.
55. Gottstein B, Hentrich B, Wyss R, Thür B, Busato A, et al. (1998) Molecular and immunodiagnostic investigations on bovine neosporosis in Switzerland. *International Journal for Parasitology* 28: 679-691.
56. Van Maanen C, Wouda W, Schares G, Von Blumröder D, Conraths FJ, et al. (2004) An interlaboratory comparison of immunohistochemistry and PCR methods for detection of *Neospora caninum* in bovine foetal tissues. *Veterinary Parasitology* 126: 351-364.
57. Baszler TV, Gay LJ, Long MT, Mathison BA (1999) Detection by PCR of *Neospora caninum* in fetal tissues from spontaneous bovine abortions. *J Clin Microbiol* 37: 4059-4064.
58. Sager H, Fischer I, Furrer K, Strasser M, Waldvogel A, et al. (2001) A Swiss case-control study to assess *Neospora caninum*-associated bovine abortions by PCR, histopathology and serology. *Veterinary Parasitology* 102: 1-15.
59. Ho MSY, Barr BC, Rowe JD, Anderson ML, Sverlow KW, et al. (1997) Detection of *Neospora* sp. from infected bovine tissues by PCR and probe hybridization. *Journal of Parasitology* 83: 508-514.
60. Peters M, Wagner F, Schares G (2000) Canine neosporosis: Clinical and pathological findings and first isolation of *Neospora caninum* in Germany. *Parasitology Research* 86: 1-7.
61. Schatzberg SJ, Haley NJ, Barr SC, deLahunta A, Olby N, et al. (2003) Use of a multiplex polymerase chain reaction assay in the antemortem diagnosis of toxoplasmosis and neosporosis in the central nervous system of cats and dogs. *American Journal of Veterinary Research* 64: 1507-1513.
62. Dubey JP, Schares G (2006) Diagnosis of bovine neosporosis. *Veterinary Parasitology* 140: 1-34.
63. Williams DJL, McGarry J, Guy F, Barber J, Trees AJ (1997) Novel ELIS A for detection of *Neospora*-specific antibodies in cattle. *Veterinary Record* 140: 328-331.
64. Björkman C, Holmdahl OJM, Uggla A (1997) An indirect enzyme-linked immunoassay (ELISA) for demonstration of antibodies to *Neospora caninum* in serum and milk of cattle. *Veterinary Parasitology* 68: 251-260.
65. Lally NC, Jenkins MC, Dubey JP (1996) Evaluation of two *Neospora caninum* recombinant antigens for use in an enzyme-linked immunosorbent assay for the diagnosis of bovine neosporosis. *Clinical and Diagnostic Laboratory Immunology* 3: 275-279.
66. Schares G, Rauser M, Söndgen P, Rehberg P, Bärwald A, et al. (2000) Use of purified tachyzoite surface antigen p38 in an ELISA to diagnose bovine neosporosis. *International Journal for Parasitology* 30: 1123-1130.
67. Jenkins MC, Caver JA, Björkman C, Anderson TC, Romand S, et al. (2000) Serological investigation of an outbreak of *Neospora caninum*-associated abortion in a dairy herd in southeastern United States. *Veterinary Parasitology* 94: 17-26.



